

SAFER CHOICES LEVEL 1 STUDENT KNOWLEDGE SURVEY

Read each question carefully and mark the answer you think is the best response.

These words are used in this survey: **infertility**, **having sex**, **STD** and **HIV**. Here,

- **infertility** means the inability to produce a child.
- **having sex** means vaginal sexual intercourse.
- **STD** stands for sexually transmitted disease or diseases. Some examples of STD are herpes, gonorrhea, syphilis, chlamydia and genital warts. Sometimes these are referred to as sexually transmitted infections (STI).
- **HIV** stands for human immunodeficiency virus.

	<i>True</i>	<i>False</i>	<i>Not sure</i>
1. Some untreated STDs can cause infertility.	<input type="radio"/> ₁	<input type="radio"/> ₂	<input type="radio"/> ₃
2. It is safe to use oils with latex condoms.	<input type="radio"/> ₁	<input type="radio"/> ₂	<input type="radio"/> ₃
3. Some STDs are curable and some are not.	<input type="radio"/> ₁	<input type="radio"/> ₂	<input type="radio"/> ₃
4. A person with an STD who looks and feels healthy cannot give the infection to others.	<input type="radio"/> ₁	<input type="radio"/> ₂	<input type="radio"/> ₃
5. Some STDs put a person at higher risk of getting infected with HIV.	<input type="radio"/> ₁	<input type="radio"/> ₂	<input type="radio"/> ₃
6. A sign of having an STD can be pain or burning when urinating (peeing).	<input type="radio"/> ₁	<input type="radio"/> ₂	<input type="radio"/> ₃
7. One reason that STDs are so common is because infected persons often do not have any signs of infection.	<input type="radio"/> ₁	<input type="radio"/> ₂	<input type="radio"/> ₃
8. The best way to use a condom is to leave some space at the tip for the sperm.	<input type="radio"/> ₁	<input type="radio"/> ₂	<input type="radio"/> ₃
9. A condom should be completely unrolled before it is placed on the penis.	<input type="radio"/> ₁	<input type="radio"/> ₂	<input type="radio"/> ₃
10. Teens must be 18 years old or older to get condoms at a store or clinic.	<input type="radio"/> ₁	<input type="radio"/> ₂	<input type="radio"/> ₃
11. About 1 out of 4 sexually active teenagers get an STD each year.	<input type="radio"/> ₁	<input type="radio"/> ₂	<input type="radio"/> ₃

12. Can the following behaviors put you at risk for getting HIV?	Yes	No	Not sure
a. Sharing needles for tattooing or piercing	<input type="radio"/> ₁	<input type="radio"/> ₂	<input type="radio"/> ₃
b. Having sex without a condom	<input type="radio"/> ₁	<input type="radio"/> ₂	<input type="radio"/> ₃
c. Donating blood	<input type="radio"/> ₁	<input type="radio"/> ₂	<input type="radio"/> ₃
d. Using the same condom twice	<input type="radio"/> ₁	<input type="radio"/> ₂	<input type="radio"/> ₃
e. Hugging	<input type="radio"/> ₁	<input type="radio"/> ₂	<input type="radio"/> ₃

13. Which of the following methods are <u>effective</u> if used correctly to protect people from STD (including HIV) and pregnancy?	<i>Protects from Pregnancy & STD/HIV</i>	<i>Protects from Pregnancy only</i>	<i>Protects from Neither</i>
a. Choosing not to have sex (abstinence)	<input type="radio"/> ₁	<input type="radio"/> ₂	<input type="radio"/> ₃
b. Using hormone based birth control (e.g., the pill, Depo-Provera shot, patch, vaginal ring)	<input type="radio"/> ₁	<input type="radio"/> ₂	<input type="radio"/> ₃
c. Using latex condoms	<input type="radio"/> ₁	<input type="radio"/> ₂	<input type="radio"/> ₃
d. Using withdrawal	<input type="radio"/> ₁	<input type="radio"/> ₂	<input type="radio"/> ₃
e. Douching (washing out the vagina)	<input type="radio"/> ₁	<input type="radio"/> ₂	<input type="radio"/> ₃

End of Knowledge Survey