

Student Name _____

Date _____

B.A.R.T. Student Knowledge Survey

Read each question carefully and mark the answer you think is the best response.

These words are used in this survey: **STD** and **HIV**. For this survey,

- **STD** stands for sexually transmitted disease or diseases. Some examples of STD are herpes, gonorrhea, syphilis, chlamydia and genital warts. Sometimes these are referred to as sexually transmitted infections (STI).
- **HIV** stands for human immunodeficiency virus.

	<i>True</i>	<i>False</i>	<i>Not sure</i>
1. Most people who have HIV look sick.	<input type="radio"/> ₁	<input type="radio"/> ₂	<input type="radio"/> ₃
2. A person can get HIV during oral sex.	<input type="radio"/> ₁	<input type="radio"/> ₂	<input type="radio"/> ₃
3. A person can get HIV in one sexual contact.	<input type="radio"/> ₁	<input type="radio"/> ₂	<input type="radio"/> ₃
4. A shower after sex reduces the risk of getting HIV.	<input type="radio"/> ₁	<input type="radio"/> ₂	<input type="radio"/> ₃
5. When people have only one sex partner, they don't need to protect themselves from HIV.	<input type="radio"/> ₁	<input type="radio"/> ₂	<input type="radio"/> ₃
6. Proper use of latex condoms helps to protect people from HIV.	<input type="radio"/> ₁	<input type="radio"/> ₂	<input type="radio"/> ₃
7. People who have HIV quickly get sick.	<input type="radio"/> ₁	<input type="radio"/> ₂	<input type="radio"/> ₃
8. By having just one sex partner at a time people can protect themselves from HIV.	<input type="radio"/> ₁	<input type="radio"/> ₂	<input type="radio"/> ₃
9. HIV doesn't go through unbroken skin.	<input type="radio"/> ₁	<input type="radio"/> ₂	<input type="radio"/> ₃
10. Semen (cum) can carry HIV.	<input type="radio"/> ₁	<input type="radio"/> ₂	<input type="radio"/> ₃

	<i>True</i>	<i>False</i>	<i>Not sure</i>
11. A person must have a lot of different sex partners to be at risk for HIV.	<input type="radio"/> ₁	<input type="radio"/> ₂	<input type="radio"/> ₃
12. People who have HIV feel quite sick.	<input type="radio"/> ₁	<input type="radio"/> ₂	<input type="radio"/> ₃
13. A person can get HIV if she shares a needle for ear piercing with someone infected with HIV.	<input type="radio"/> ₁	<input type="radio"/> ₂	<input type="radio"/> ₃
14. People who practice withdrawal during sex won't get HIV.	<input type="radio"/> ₁	<input type="radio"/> ₂	<input type="radio"/> ₃
15. A negative result on the HIV test can happen even if somebody has HIV.	<input type="radio"/> ₁	<input type="radio"/> ₂	<input type="radio"/> ₃
16. Breast milk can carry HIV.	<input type="radio"/> ₁	<input type="radio"/> ₂	<input type="radio"/> ₃
17. Most people who have HIV know they have it.	<input type="radio"/> ₁	<input type="radio"/> ₂	<input type="radio"/> ₃
18. No case of AIDS was ever caused by kissing on the cheek.	<input type="radio"/> ₁	<input type="radio"/> ₂	<input type="radio"/> ₃
19. All STDs can be cured.	<input type="radio"/> ₁	<input type="radio"/> ₂	<input type="radio"/> ₃
20. Infection with other STDs can put a person at higher risk of getting HIV during sex.	<input type="radio"/> ₁	<input type="radio"/> ₂	<input type="radio"/> ₃

End of Knowledge Survey